

Aminogro[®]

Combi

Fe 5% Mn 2% Zn 4%
Cu 1% B 0.6% Mo 0.07%

MAXi

Organic Carbon: 16% w/w Amine Nitrogen: 7.4% w/w

organaPLEX L-form Amino Acid
Chelate Carboxylate Mannitol Complex

PLANT FUNCTION

- Cell wall strength
- Sugar transportation
- Flower retention, pollen formation, fruit set
- Seed germination
- Chlorophyll production
- Hormone and enzyme synthesis



ORGANIC CROP PROTECTANTS PTY LTD
61 Turrella St, Turrella NSW 2205 Australia
Telephone: 1800 634 204



CONTENTS
10 kg

GENERAL INFORMATION

Aminogro MAXi OrganoPLEX provides plants with COMBI micronutrient chelates using L-form free amino acids and other simple organic sugars and acids. Nutrients are absorbed and shuttled around inside plants in the form of amino acids, simple sugars and acids so OrganoPLEX is an efficient way of delivering and mobilizing these micronutrients in plants. The organic carbon in Aminogro MAXi also enhances microbial activity in soil which assists the plant accessing COMBI micronutrients when applied to the soil.

DIRECTIONS FOR USE: 1 L Scoop = approx. 300 g

The rates below are a guide and dependent upon soil and sap tests throughout the growing season. Consult your OCP Agronomist for further advice.

Tree Crop	Rate	Application Timing	Field Crops	Rate	Application Timing
Pome Fruit	2.5-3 kg/ha	1. At fruit set 2. One month after fruit set 3. 2 months before harvest 4. 2 weeks before harvest	Lettuce	2 kg/ha	1. 4 leaf stage 2. Every 14 days until harvest
Stone Fruit	1.5-2 kg/ha	1. At fruit set 2. 2-3 weeks later 3. 2-3 weeks later 4. 2 weeks before harvest	Carrots, Onions, Leeks, Turnips	2 kg/ha	1. 2 weeks after emergence 2. At root enlargement 3. 6 weeks before harvest
Nuts	1-2 kg/ha	1. At bud swell 2. At nut set 3. 2 weeks later	Brassica, Crucifer	2 kg/ha	1. 6-8 true leaf stage 2. 10-14 days later 3. At head initiation
Bananas	2-3 kg/ha	With seasonal application of Potassium and Nitrogen	Bean & peas	2 kg/ha	1. 4-6 true leaf stage 2. At first bloom 3. At first pod
Vine Crop	Rate	Application Timing	Potatoes, Yam, Taro	2 kg/ha	1. (Seed treatment optional) 2. At tuber set 3. Early bloom
Table Grapes	2.5-3 kg/ha	1. Pre-bloom E-L 12 2. Pre-bloom E-L 16 3. Fruit set E-L 27 4. Post harvest E-L	Sweet corn	2 kg/ha	1. 2-6 true leaf stage 2. 50-75cm growth 3. Just prior to tasselling
Wine Grapes	2 kg/ha	1. Pre-bloom E-L 12 2. Fruit set E-L 27 3. Post harvest E-L 40	Protected Crops	Rate	Application Timing
Strawberries/ Blueberries	2-2.5 kg/ha	Begin at fruit set and repeat as sap results indicate	Hydroponic NFT All crops	500 g -1 kg/ 10,000 L	Consult OCP agronomist
			Nursery Stock	Foliar	If deficiency is detected consult OCP agronomist

Professional Turf	Rate	Application Timing	General Information
Greens, Tee & intensively managed Sports turf & Fairways	5 kg/ha	Monthly applications or as required to assist with wear and disease resistance	Aminogro MAXi COMBI is compatible with most fungicides and insecticides. If mixing in alkaline water, adjust to pH 6-6.5 using a suitable buffer

COMPATIBILITY

Aminogro MAXi can be mixed with most horticultural sprays. However, the performance of some sprays may be affected by high Cation levels like Calcium, Potassium and Phosphorus. It is best to apply these sprays separately or conduct a jar test to determine compatibility in the ratios they are to be used. Always check pesticide labels for compatibility with foliar fertilizers, or test the mixture on a small area before widespread application.

BATCH NO.
DATE OF MANUFACTURE