

208

50

152

# Aminogro<sup>®</sup>

## 12% Copper

### MAXi

Organic Carbon: 18% w/w Amine Nitrogen: 6% w/w

**organaPLEX** L-form Amino Acid  
Chelate Carboxylate Mannitol Complex

#### PLANT FUNCTION

- Photosynthesis
- Enzyme production
- Carbohydrate & nitrogen metabolism
- Lignin synthesis for cell wall strength and root cap

#### DEFICIENCY SYMPTOMS

- Dieback of stems & twigs
- Yellowing of leaves
- Stunted growth



ORGANIC CROP PROTECTANTS PTY LTD  
61 Turrella St, Turrella NSW 2205 Australia  
Telephone: 1800 634 204



CONTENTS  
**10 kg**

#### GENERAL INFORMATION

Aminogro MAXi OrganoPLEX provides plants with a Copper chelate using L-form free amino acids and other simple organic sugars and acids. Nutrients are absorbed and shuttled around inside plants in the form of amino acids, simple sugars and acids so OrganoPLEX is an efficient way of delivering and mobilizing Copper in plants. The organic carbon in Aminogro MAXi also enhances microbial activity in soil which assists the plant accessing Copper and other nutrients when applied to the soil.

#### DIRECTIONS FOR USE: 1 L Scoop = approx. 250 g

The rates below are a guide and dependent upon soil and sap tests throughout the growing season. Consult your OCP Agronomist for further advice.

Tree Crop	Rate	Application Timing	Field Crops	Rate	Application Timing
Pome Fruit	800 g-1.5 kg/ha	1-2 weeks before flowering	Cereal, Soybeans	1 kg/ha	As required by soil test results
Stone Fruit	800 g-1.5 kg/ha	At first flush	Leafy veggies, Lettuce	500 g/ha	4 leaf stage
Nuts	800 g-1.5 kg/ha	At first flush	Carrots, Onions, Leeks, Turnips	500 g/ha	2 weeks after emergence
Bananas	500 g-1 kg/ha	With seasonal application of Potassium and Nitrogen	Brassica, Crucifer	500 g/ha	6-8 true leaf stage
Vine Crop	Rate	Application Timing	Bean & peas	500 g/ha	4-6 true leaf stage
Table Grapes	500 g-1.5 kg/ha	2 weeks before flowering	Potatoes, Yam, Taro	500 g/ha	At tuber set
Wine Grapes	800 g-1.5 kg/ha	2 weeks before flowering	Sweet corn	500 g/ha	2-6 true leaf stage
Strawberries/Blueberries	500 g-1 kg/ha	Begin just before flowering & repeat as sap results indicate	Tomato, Capsicum	500-750 g/ha	15-20cm
Protected Crops	Rate	Application Timing	Cucurbits	500 g/ha	4 true leaf
Hydroponic run to waste & NFT crops	100 g/10,000 L	Consult OCP agronomist for crop specific requirements	Professional Turf	Rate	Application Timing
Nursery Stock	Foliar	If deficiency is detected consult OCP agronomist	Greens & Tees	100 g/1000 m <sup>2</sup>	As required by soil test. Higher rates may be used during time of disease pressure
			Fairways	500 g/ha	As required by soil test

#### COMPATIBILITY

Aminogro MAXi can be mixed with most horticultural sprays. However, the performance of some sprays may be affected by high Cation levels like Calcium, Potassium and Phosphorus. It is best to apply these sprays separately or conduct a jar test to determine compatibility in the ratios they are to be used. Always check pesticide labels for compatibility with foliar fertilizers, or test the mixture on a small area before widespread application.

BATCH NO.  
DATE OF MANUFACTURE

170