

208

50

152

Aminogro[®]

N13

MAXi

Organic Nitrogen 13% w/w

Aminogro MAXi N13 provides plants with an array of amino acids in the form of L-form free amino acids, peptides and polypeptides. Nitrogen is carried around inside plants in the form of amino acids so fertilizing with amino acids is an efficient way of delivering Nitrogen to the plant. The organic carbon in Aminogro MAXi N13 also enhances microbial activity which assists the plant to access other nutrients. Aminogro MAXi N13 also acts as a natural chelator of other nutrients in tank mixes.



ORGANIC CROP PROTECTANTS PTY LTD
61 Turrella St, Turrella NSW 2205 Australia
Telephone: 1800 634 204



CONTENTS
10 kg

DIRECTIONS FOR USE:

The rates below are a guide and dependent upon the amount of Nitrogen applied as a basal fertiliser pre-season and crop yield targets. Adjustments should be made to Nitrogen requirements throughout the growing season based on sap tests. Consult your OCP Agronomist for further advice.

Tree Crop	Rate	Application timing	Field Crops	Rate	Application timing
Pome Fruit	500 g/100 L or max 5 kg/ha	1. Green tip to tight cluster 2. Open cluster to pink 3. 1st cover to 4th cover 4. Every 2-3 wks after 1st cover 5. 2-3 wks before harvest	Cereal, Soybeans	2-3 kg/ha	As required by soil or sap test results.
Stone Fruit	500 g/100 L or max 5 kg/ha	1. At petal fall 2. 10-14 days later 3. Post harvest-drip x 2 apps 10kg/ha	Lettuce	500 g/100 L or max 10 kg/ha	1. Seed treatment 2. 4 leaf stage 3. every 14 days until harvest
Nuts	500 g/100 L or max 5 kg/ha	1. At petal fall 2. 10-14 days later 3. Post harvest-drip x 2 apps 10kg/ha	Carrots, Onions, Leeks, Turnips	500 g/100 L or max 10 kg/ha	1. 2 weeks after emergence 2. At root enlargement 3. 10-14 days until harvest
Bananas	1 kg/100 L or max 50 kg/ha	With every spray or as required through fertigation	Brassica, Crucifer	500 g/100 L or max 10 kg/ha	1. 4-6 true leaf stage 2. 10-14 days later 3. At head initiation
Vine crop	Rate	Application timing	Bean & peas	500 g/100 L or max 10 kg/ha	1. 4-6 true leaf stage 2. At first bloom 3. At first pod
Table Grapes	500 g/100 L or max 5 kg/ha	1. 10-15 cm canes 2. 25-35 cm canes 3. Berry set 4. 7 days later 5. 7 days later Post harvest-drip x 2 apps	Potatoes, Yam, Taro	500 g/100 L or max 10 kg/ha	1. (Seed treatment optional) 2. At tuber set 3. 10-14 days later 4. Early bloom
Wine Grapes	300 g/100 L or max 3 kg/ha	1. Berry set 2. Veraison* 3. Pre-bloom* 4. Berry set* 5. Veraison* Post harvest - drip x 2 apps 10kg/ha * For growers who want bigger berries and better set	Sweet corn	500 g/100 L or max 10 kg/ha	1. 2-6 true leaf stage 2. 50-75cm growth 3. Just prior to tasselling
Strawberries/ Blueberries	5 kg/ha	Begin at fruit set and repeat as sap results indicate	Protected crops	Rate	Application timing
			Hydroponic NFT All crops	5 kg/10,000 L	Every two weeks. Consult OCP agronomist if using the product for entire Nitrogen source
			Cutflowers	Drip & Foliar	Every 2nd wk @ 2.5kg / 2000 plants as a drip and 250 g / 100L as foliar every wk
			Nursery Stock	Foliar	Every week @ 500g / 100L

Professional turf	Rate	Application timing	General Information
Greens, Tee & intensively managed Sports turf	5 kg/ha	Monthly applications or as required for aesthetic appeal	Aminogro MAXi Iron is compatible with PGR's. If mixing in alkaline water, adjust to pH 6-6.5 using a suitable buffer
Fairways & Sports Fields	2 kg/ha	Monthly application for optimum greening	

COMPATIBILITY

Aminogro MAXi can be mixed with most horticultural sprays. Always check pesticide labels for compatibility with foliar fertilizers, or test the mixture on a small area before widespread application.

BATCH NO.
DATE OF MANUFACTURE