

208

50

152

Aminogro[®]

N7

MAXi

Organic Nitrogen 7% w/v

Aminogro MAXi N7 provides plants with an array of amino acids in the form of L-form free amino acids, peptides and polypeptides. Nitrogen is carried around inside plants in the form of amino acids so fertilizing with amino acids is an efficient way of delivering Nitrogen to the plant. The organic carbon in Aminogro MAXi N7 also enhances microbial activity which assists the plant to access other nutrients. Aminogro MAXi N7 also acts as a natural chelator of other nutrients in tank mixes.



ORGANIC CROP PROTECTANTS PTY LTD
61 Turrella St, Turrella NSW 2205 Australia
Telephone: 1800 634 204



CONTENTS
20 L

DIRECTIONS FOR USE:

The rates below are a guide and dependent upon the amount of Nitrogen applied as a basal fertiliser pre-season and crop yield targets. Adjustments should be made to Nitrogen requirements throughout the growing season based on sap tests. Consult your OCP Agronomist for further advice.

Tree Crop	Rate	Application timing	Field Crops	Rate	Application timing
Pome Fruit	1 L/100 L or max 10 L/ha	1. Green tip to tight cluster 2. Open cluster to pink 3. 1st cover to 4th cover 4. Every 2-3 wks after 1st cover 5. 2-3 wks before harvest	Cereal, Soybeans	max 10 L/ha 1 L/100 L or	As required by soil or sap test results.
Stone Fruit	1 L/100 L or max 10 L/ha	1. At petal fall 2. 10-14 days later 3. Post harvest-drip x 2 apps 20L/ha	Lettuce	max 10 L/ha	1. Seed treatment 2. 4 leaf stage 3. every 14 days until harvest
Nuts	1 L/100 L or max 10 L/ha	1. At petal fall 2. 10-14 days later 3. Post harvest-drip x 2 apps 20L/ha	Carrots, Onions, Leeks, Turnips	1 L/100 L or max 10 L/ha	1. 2 weeks after emergence 2. At root enlargement 3. 10-14 days until harvest
Bananas	1 L/100 L or max 100 L/ha	With every spray or as required through fertigation	Brassica, Crucifer	1 L/100 L or max 10 L/ha	1. 4-6 true leaf stage 2. 10-14 days later 3. At head initiation
Vine crop	Rate	Application timing	Bean & peas	1 L/100 L or max 10 L/ha	1. 4-6 true leaf stage 2. At first bloom 3. At first pod
Table Grapes	1 L/100 L or max 100 L/ha	1. 10-15 cm canes 2. 25-35 cm canes 3. Berry set 4. 7 days later 5. 7 days later Post harvest-drip x 2 apps	Potatoes, Yam, Taro	1 L/100 L or max 10 L/ha	1. (Seed treatment optional) 2. At tuber set 3. 10-14 days later 4. Early bloom
Wine Grapes	500 ml/100 L or max 5 L/ha	1. 2-5 cm bunch 2. 14 days later (at least 14 days prior to first bloom) 3. Pre-bloom* 4. Berry set* 5. Veraison* Post harvest - drip x 2 apps 20L/ha* * For growers who want bigger berries and better set	Sweet corn	1 L/100 L or max 10 L/ha	1. 2-6 true leaf stage 2. 50-75cm growth 3. Just prior to tasselling
Strawberries & Blueberries	10 L-20 L/ha	Begin at fruit set and repeat as sap results indicate	Protected crops	Rate	Application timing
			Hydroponic NFT All crops	2-5 L/10,000 L	Every two weeks. Consult OCP agronomist if using the product for entire Nitrogen source
			Cutflowers	Drip & Foliar	Every 2nd wk @ 1L / 2000 plants as a drip and 200 ml / 100L as foliar every wk
			Nursery Stock	Foliar	Every week @ 100ml / 100L
			Professional turf	Rate	Application Timing
			Greens & Tees	500 mL/100 m ²	Renovation As required by soil test and growing conditions
			Greens & Tees	250 ml/100 m ²	Maintenance As required by soil test and growing conditions
			Fairways	20 L/ha	As required by soil test and growing conditions

COMPATIBILITY

Aminogro MAXi can be mixed with most horticultural sprays. Always check pesticide labels for compatibility with foliar fertilizers, or test the mixture on a small area before widespread application.

BATCH NO.
DATE OF MANUFACTURE