

8.5cm

CAUTION
 KEEP OUT OF REACH OF CHILDREN
 READ SAFETY DIRECTIONS BEFORE OPENING OR USING

AzaMax[®]

INSECTICIDE

ACTIVE CONSTITUENTS: 11.82g/L AZADIRACTIN A and B
 present as 29.55 g/L *Azadirachta indica* extract

GROUP **22A** INSECTICIDE

For the control of two-spotted mites, aphids and whitefly in
 Floriculture and ornamentals and fungus gnats in potting
 soil as shown in the Directions for Use.



Mites



Aphids



Whitefly



Fungus Gnats



ORGANIC CROP PROTECTANTS Pty Ltd
 42 Halloran Street, Lilyfield NSW 2040
 Telephone (02) 9810 4566 Fax (02) 9810 4674



STORAGE AND DISPOSAL: Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS: Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing product for use, wear elbow length PVC gloves. Wash hands after use. After each day's use wash gloves.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

MSDS: Additional information is listed in the Material Safety Data Sheet.

NOTE:

Because Organic Crop Protectants Pty Ltd has no control over handling, storage or use of AzaMax Insecticide, the company makes no express or implied representations or warranty (other than non-excludable statutory warranties). This product is sold on the condition that the buyer and/or user assumes all risks of the handling, use of AzaMax Insecticide and results obtained from its use.

For further information contact 1800 634 204
 8am-5pm Monday to Friday. ® Registered trademark

Manufactured by: EID Parry (India) Limited



DIRECTIONS FOR USE

RESTRAINT: DO NOT use on plants that produce food for human or animal consumption.

Crop/ Situation	Pest	Rate	Critical Comments
Floriculture, Ornamentals	Two-spotted Mites Whitefly Aphids	200 mL to 300 mL per 100 L water	Regularly monitor plants for presence of pests. Use a 7 to 10 day spray regime when pests are present. Multiple applications (3-5) may be required. Use the higher rate under higher pest pressure. Spray volume will vary according to the crop and location. Apply in a spray volume of 250 to 500 L/ha, using a hand held wand or knapsack spray. Addition of a botanical oil concentrate adjuvant is recommended. Note plant safety PRECAUTION below.
Potting soil (for floriculture and ornamentals only)	Fungus gnats	30 mL/10 L water	Apply as a soil drench using approximately 1 litre mixture per 8 litres potting soil.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD (WHP): NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS: AzaMax is a biologically derived product that is active on the above sucking insect pests on a wide range of ornamental plants. Consistent applications are necessary during pest activity.

PRECAUTION - Plant Safety: Trials on sensitive plants such as ferns, poinsettias and African violets have resulted in crop damage. Minor phytotoxic effects have occurred in other species. Testing in a small area is therefore recommended before spraying to determine sensitive species and varieties.

MIXING: Half fill spray tank and commence agitation. Slowly add required amount of AzaMax. Top up with required quantity of water. Agitate well before commencing spray procedure.

INSECTICIDE RESISTANCE WARNING GROUP **22A** INSECTICIDE

For insecticide resistance management AzaMax Insecticide is a Group 22A insecticide. Some naturally occurring insect biotypes resistant to AzaMax Insecticide and other Group 22A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if AzaMax Insecticide or other Group 22A insecticides are used repeatedly. The effectiveness of AzaMax Insecticide on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Organic Crop Protectants Pty Ltd accepts no liability for any losses that may result from the failure of AzaMax Insecticide to control resistant insects. AzaMax Insecticide may be subject to specific resistance management strategies. For further information contact your local supplier, Organic Crop Protectants Pty Ltd representative or local agricultural department agronomist.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT: Dangerous to aquatic invertebrates. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

APVMA Approval Number 61980/1L/0809
 DOM EXPIRY BATCH NO.

31.0314cm