

CAUTION

KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING

INSECTICIDE

ACTIVE CONSTITUENTS: 11.82 g/L AZADIRACHTIN A and B present as 29.55 g/L Azadirachta indica extract

GROUP **INSECTICIDE**

For the control of two-spotted mites, aphids and whitefly in floriculture and ornamentals and fungus gnats in potting soil as shown in the Direction for Use.





Aphids





Process

Flexo Lytho Gravure

Offset

Whitefly



CONTENTS 5 L

ORGANIC CROP PROTECTANTS 1956 Dandenong Rd, Clayton VIC 3168 Telephone: 1800 634 204 (Free Call)





Job OCP658_Azamax Label_5L

Version

Size 140x135 mm









Barcode % 100





PLEASE NOTE. Redpin strives to ensure accuracy of final artwork. However, the responsibility for final sign-off rests with the client. Redpin cannot accept responsibility for incorrect client approved artwork.

94 Harris Street, Harris Park NSW 2150 T. 02 9683 1177 E. creative@redpin.com.au W. redpin.com.au

DIRECTIONS FOR USE

RESTRAINT: DO NOT use on plants that produce food for human or animal consumption.

CROP/SITUATION	PEST	RATE	CRITICAL COMMENTS
Floriculture, Ornamentals	Two-spotted Mites Whitefly Aphids	200 mL to 300 mL per 100L water	Regularly monitor plants for presence of pests. Use a 7 to 10 day spray regime when pests are present. Multiple applications (3-5) may be required. Use the higher rate under higher pest pressure. Spray volume will vary according to the crop and location. Apply in a spray volume of 250 to 500 L/ha, using a hand held wand or knapsack spray. Addition of a botanical oil concentrate adjuvant is recommended. Note plant safety PRECAUTION below.
Potting soil (for Floriculture and Ornamentals only)	Fungus Gnats	30 mL/ 10L water	Apply as a soil drench using approximately 1 litre mixture per 8 litres potting soil.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD (WHP): NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS: AzaMax® is a biologically derived product that is active on the above sucking insect pests on a wide range of omamental plants. Consistent applications are necessary during pest activity.

STORAGE AND DISPOSAL: Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank.

MIXING: Half fill spray tank and commence agitation. Slowly add required amount of AzaMax®. Top up with required quantity of water. Agitate well before

INSECTICIDE RESISTANCE WARNING

BATCH NO:

DOM:

GROUP UN INSECTICIDE

For insecticide resistance management AzaMax® Insecticide is a Group UN insecticide. Some naturally occurring insect biotypes resistant to AzaMax® Insecticide and other Group UN insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if AzaMax® Insecticide or other Group UN insecticides are used repeatedly. The effectiveness of AzaMax® Insecticide on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Organic Crop Protectants accepts no liability for any losses that may result from the failure of AzaMax® Insecticide to control resistant insects.

AzaMax® Insecticide may be subject to specific resistance management strategies. For further information contact your local supplier, Organic Crop Protectants representative or local agricultural department agronomist.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT: Dangerous to aquatic invertebrates. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

AzaMax is a registered trade mark of Coromandel nternational and Organic Crop Protectants The OCP drop device is a registered trade mark of

APVMA Approval Number 61980/5L/0809 Manufactured by: Coromandel International Contains NeemAzal by COROMANDEL

PRECAUTION - PLANT SAFETY: Trials on sensitive plants such as ferns, poinsettias and African violets have resulted in crop damage. Minor phytotoxic effects have occurred in other species. Testing in a small area is therefore recommended before spraying to determine sensitive species and varieties. marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt. SAFETY DIRECTIONS: Will irritate the eyes and skin. Avoid contact with eyes and

skin. When opening the container and preparing product for use, wear elbow length PVC gloves. Wash hands after use. After each day's use wash gloves. FIRST AID: If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone Australia 131 126). If medical advice is needed, have product container or label at hand.

SDS: Additional information is listed in the Safety Data Sheet available from the

FOR EMERGENCIES ONLY, CALL AUSTRALIA 1800 220 770 (ALL HOURS)



100% Magnification Barcode 9336099555734