




Organic Crop Protectants

ecoprotector™

FUNGICIDAL POTASSIUM SOAP

ACTIVE CONSTITUENTS: 194 g/L POTASSIUM SALTS OF FATTY ACIDS

FOR THE CONTROL OF BOTRYTIS IN
Grapes



CONTENTS
20L

ORGANIC CROP PROTECTANTS
1956 Dandenong Rd, Clayton VIC 3168
Telephone: 1800 634 204 (Free Call)
ocp.com.au

photos courtesy of PIRSA-SARDI



Restricted for use under ACO Standards

DIRECTIONS FOR USE

CROP	DISEASE	RATE	CRITICAL COMMENTS
Grapevines	Botrytis	20 litres per ha in 500 to 1000 litres water	Apply by high volume spraying to the point of run-off. Ensure thorough coverage of all parts of treated plants. Higher water rates will give better coverage and better botrytis control. Apply the first treatment at early flowering followed by a second application around late flowering. Repeat applications at other strategic growth stages e.g. 20% Capfall, 80% Capfall and Bunch Closing or at 21 day intervals until 14 days before harvest.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD (WHP):

GRAPEVINES: DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.

GENERAL INSTRUCTIONS

ecoprotector™ does not control either powdery or downy mildew. ecoprotector™ can be mixed with sulfur for powdery mildew control. A tank mixture with a copper hydroxide for the control of downy mildew also gives good control of powdery mildew when applied as a preventative spray. Critical times for botrytis control are during flowering, pre-bunch closure, veraison and the period up to harvest when the berries are maturing. If ecoprotector™ is applied only at these times control of botrytis will be obtained but the level of control will not be as high as that achieved with a regular program of application at 21 day intervals.

MIXING: Add the product at the appropriate rate to water and mix well. Use good quality, soft water of pH 6 to 8. The use of hard water may cause precipitation and the addition of a softener is recommended.

APPLICATION: ecoprotector™ should only be applied using suitable dilute ground application equipment. Concentrate spraying is not recommended. Equipment should produce a spray that ensures penetration of the canopy and coverage of the flowers or bunches. Use a sprayer designed to apply high spray volumes, up to the point of run-off and matched to the crop being sprayed. The sprayer should be set-up and operated to achieve even coverage throughout the crop canopy to the point of run-off. Avoid excessive run-off. Add the amount of product specified in the Directions for Use table. However, if greater than 1000 L/ha is required to achieve point of run-off, adjust the amount of product by 2 L of ecoprotector™ for every 100 L to ensure that a dilution of 2% is achieved. The required dilute spray volume to achieve point of run-off will change and the sprayer set up and operation may also need to be changed, as the crop grows.

COMPATIBILITY: ecoprotector™ is compatible with copper hydroxide fungicides and sulfur. Do not add other pesticides. The addition of wetting or sticking agents is not recommended.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT: DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL: KEEP OUT OF REACH OF CHILDREN. DO NOT store at low temperatures (below 5°C). Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

SDS: Additional information is listed in the Safety Data Sheet available from the supplier and ocp.com.au.

For further information contact Organic Crop Protectants

Free Call: 1800 634 204 Email: info@ocp.com.au

FOR EMERGENCIES ONLY, CALL AUSTRALIA 1800 033 111 (ALL HOURS)

ecoprotector and the OCP drop device are trade marks of Organic Crop Protectants.

APVMA Approval No. 63223 / 118687

BATCH NO.
DATE OF MANUFACTURE

9336099556298
100% MAGNIFICATION