

208

50

152

Aminogro[®]

12% Iron

MAXi

Organic Carbon: 19% w/w Amine Nitrogen: 7.6% w/w

organaPLEX L-form Amino Acid
Chelate Carboxylate Mannitol Complex

PLANT FUNCTION

- Chlorophyll production
- Photosynthesis
- Nitrogen uptake & lignin formation

DEFICIENCY SYMPTOMS

- Yellowing of leaves and interveinal chlorosis (green veins and yellowing between veins)



ORGANIC CROP PROTECTANTS PTY LTD
61 Turrella St, Turrella NSW 2205 Australia
Telephone: 1800 634 204



CONTENTS
10 kg

GENERAL INFORMATION

Aminogro MAXi OrganoPLEX provides plants with an Iron chelate using L-form free amino acids and other simple organic sugars and acids. Nutrients are absorbed and shuttled around inside plants in the form of amino acids, simple sugars and acids so OrganoPLEX is an efficient way of delivering and mobilizing Iron in plants. The organic carbon in Aminogro MAXi also enhances microbial activity in soil which assists the plant accessing Iron and other nutrients when applied to the soil.

DIRECTIONS FOR USE: 1 L Scoop = approx. 400 g

The rates below are a guide and dependent upon soil and sap tests throughout the growing season. Consult your OCP Agronomist for further advice.

Tree Crop	Rate	Application Timing	Field Crops	Rate	Application Timing
Pome Fruit	1-2 kg/ha	2 weeks after fruit set	Cereal, Soybeans	2-3 kg/ha	As required by soil test results
Stone Fruit	1-2 kg/ha	1-2 weeks after fruit set	Leafy veggies	1-2 kg/ha	4 leaf stage
Nuts	1-2 kg/ha	2-3 weeks after nut set	Lettuce		
Vine Crop	Rate	Application Timing	Carrots, Onions, Leeks, Turnips	2 kg/ha	2 weeks after emergence
Table Grapes	1-1.5 kg/ha	Pre-bloom E-L 12 & as required by sap & soil test	Brassica, Crucifer	2 kg/ha	6-8 true leaf stage
Wine Grapes	750 g-1.5 kg/ha	Pre-bloom E-L 12 & as required by sap test	Bean & peas	2 kg/ha	4-6 true leaf stage
Strawberries/ Blueberries	750 g-1.5 kg/ha	Before flowering & as required by sap & soil test	Potatoes, Yam, Taro	2 kg/ha	At tuber set
Protected Crops	Rate	Application Timing	Sweet corn	2 kg/ha	1. 2-6 true leaf stage 2. 50-75cm growth
Hydroponic NFT All crops	200 g/10,000 L	Consult OCP agronomist	Cucurbits	2 kg/ha	4 true leaf stage
Nursery Stock	Foliar	If deficiency is detected consult OCP agronomist	Tomato, Capsicum	750 g-1 kg/ha	15-20cm
Professional Turf	Rate	Application Timing	Greens & Tees	500 g/1000 m ²	As required for colour
			Fairways	2 kg/ha	As required for colour

COMPATIBILITY

Aminogro MAXi can be mixed with most horticultural sprays. However, the performance of some sprays may be affected by high Cation levels like Calcium, Potassium and Phosphorus. It is best to apply these sprays separately or conduct a jar test to determine compatibility in the ratios they are to be used. Always check pesticide labels for compatibility with foliar fertilizers, or test the mixture on a small area before widespread application.

BATCH NO.
DATE OF MANUFACTURE

170