

CAUTION

KEEP OUT OF REACH OF CHILDREN

SPLAT CL-NR

200g/kg Cue-lure parapheromone for the attraction of
Male Queensland fruit fly (*Bactrocera tryoni*)

SPLAT (Specialized Pheromone and Lure Application Technology) is an amorphous polymer matrix for the sustained controlled release of insect pheromones. SPLAT CL is used as part of the ANAMED protein baiting program for Female QLD Fruit Fly.

Distributed by: ORGANIC CROP PROTECTANTS

1956 Dandenong Road, Clayton VIC 3168

Freecall: 1800 6342 04 E: info@ocp.com.au

NET CONTENTS 15kg



HOW TO USE

RATE: 300g/ha as small 3g dollops delivering 100 dollops/ha evenly throughout the crop. Add the required amount of toxicant as per instructions provided by manufacturers label.

TIMING: Apply every 4-8 weeks at first signs of fruit fly activity via monitoring with ISCA Ball traps or similar. Apply in combination with ANAMED female fruit fly baiting program. Frequency of application depends on pest pressure, geographic area, time of year and toxicant's field life.

MIXING & DISPENSING: Refer to manufacturers instructions for mixing toxicant into baits. Mix toxicant until thoroughly dispersed into the SPLAT matrix. To determine a 3g dollop wear rubber or latex gloves and using a tongue depressor or ice cream stick dispense a teaspoon size of quantity of SPLAT CL onto a weighing scale with an accuracy of 0.1g until 3g is reached.

APPLICATION: Apply the dollops evenly to the boundary trees first at approx. 10m spacing and then work your way through the paddock at approx. 10m intervals. Depending on type of crop apply the dollops to stakes, non-fruiting branches/tree trunks (center of canopy), fences, non edible border vegetation of the surrounding crop fields. Make an effort to apply to less environmentally exposed or high traffic areas to minimize dislodgement.

FIRST AID STATEMENT

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Wash skin immediately with soap and water, then rinse with plenty of water for 15-20 minutes.

FIRST AID MEASURES

If poisoning occurs, contact a doctor or **Poisons Information Centre (Phone Australia 131 126)**. If any enquires regarding misuse of product, emergencies or enquiries relating to pets/humans ingesting product, please direct them to our Emergency Response Service (**ERS**) In an emergency, call **1800 033 111 (ALL HOURS)**

Batch Number:

