

208



Aminogro[®]

12% Iron

MAXi

Organic Carbon: 19% w/w Amine Nitrogen: 7.6% w/w

OrganoPLEX[®] L-form Amino Acid
Chelate Carboxylate Mannitol Complex

PLANT FUNCTION

- Chlorophyll production
- Photosynthesis
- Nitrogen uptake & lignin formation

DEFICIENCY SYMPTOMS

- Yellowing of leaves and interveinal chlorosis (green veins and yellowing between veins)



ORGANIC CROP PROTECTANTS
1956 Dandenong Rd, Clayton VIC 3168
Telephone: 1800 634 204 (Free Call)
ocp.com.au



CONTENTS
25 kg

Y20-3316

208

GENERAL INFORMATION

Aminogro[®] MAXi Iron provides plants with an iron chelate using L-form free amino acids and other simple organic sugars and acids. Nutrients are absorbed and shuttled around inside plants in the form of amino acids, simple sugars and acids so OrganoPLEX[™] is an efficient way of delivering and mobilising iron in plants. The organic carbon in Aminogro[®] MAXi Iron also enhances microbial activity in soil which assists the plant accessing iron and other nutrients when applied to the soil.

DIRECTIONS FOR USE:

The rates below are a guide and dependent upon soil and sap tests throughout the growing season. Consult your OCP Agronomist for further advice.

Tree Crop	Rate	Application Timing
Pome Fruit	1-2 kg/ha	2 weeks after fruit set
Stone Fruit	1-2 kg/ha	1-2 weeks after fruit set
Nuts	1-2 kg/ha	2-3 weeks after nut set
Vine Crop	Rate	Application Timing
Table Grapes	1-1.5 kg/ha	Pre-bloom E-L 12 & as required by sap & soil test
Wine Grapes	750 g-1.5 kg/ha	Pre-bloom E-L 12 & as required by sap test
Strawberries/Blueberries	750 g-1.5 kg/ha	Before flowering & as required by sap & soil test
Protected Crops	Rate	Application Timing
Hydroponic NFT All Crops	200 g/10,000 L	Consult OCP agronomist
Nursery Stock	Foliar	If deficiency is detected consult OCP agronomist

Field Crops	Rate	Application Timing
Cereal, Soybeans	2-3 kg/ha	As required by soil test results
Leafy Veggies Lettuce	1-2 kg/ha	4 leaf stage
Carrots, Onions, Leeks, Turnips	2 kg/ha	2 weeks after emergence
Brassica, Crucifer	2 kg/ha	6-8 true leaf stage
Bean & Peas	2 kg/ha	4-6 true leaf stage
Potatoes, Yam, Taro	2 kg/ha	At tuber set
Sweet Corn	2 kg/ha	1. 2-6 true leaf stage 2. 50-75cm growth
Cucurbits	2 kg/ha	4 true leaf stage
Tomato, Capsicum	750 g-1 kg/ha	15-20cm
Professional Turf	Rate	Application Timing
Greens & Tees	500 g/1000 m ²	As required for colour
Fairways	2 kg/ha	As required for colour

COMPATIBILITY

Aminogro[®] MAXi Iron can be mixed with most horticultural sprays. However, the performance of some sprays may be affected by high cation levels like calcium, potassium and phosphorus. It is best to apply these sprays separately or conduct a jar test to determine compatibility in the ratios they are to be used. Always check pesticide labels for compatibility with foliar fertilisers, or test the mixture on a small area before widespread application.

For further information contact Organic Crop Protectants
Free Call: 1800 634 204 Email: info@ocp.com.au
FOR EMERGENCIES ONLY, CALL AUSTRALIA 1800 033 111 (ALL HOURS)

Aminogro and the OrganoPLEX device are registered trade marks and the OCP drop device is a trade mark of Organic Crop Protectants



Y20-3322



Job OCP572 AminogroMaxi IRON 25kg

Version

Size 208mm x 190mm
Colours



Barcode % 100%
Spot



Process
Flexo
Lytho
Gravure
Offset

FA

04/05/20