



The ecocopper Advantage

- ✓ **Optimised particle** size distribution and shape for greater leaf coverage and contact with bacteria and fungi
- ✓ **Balanced release** of (fast & slow) copper ions and released when needed due to humidity & CO₂ reaction.
- ✓ **Longer persistence** than other copper based fungicides
- ✓ **Greater copper stability** under wider pH range – lower phytotoxicity to the crop due to less “dumping” of copper
- ✓ **Excellent dispersion** and suspensibility due to fluid bed WG formulating technology
- ✓ **Input for Organic Production** – Registered for use in organic farming systems. ACO & NOP Approved
- ✓ **Instantaneous Dispersion** Dissolves completely in water, with little to no residue left in the spray tank (*see images right*)

Super fine granules for easy dispersion in water



ecocopper™

375 WG ORGANIC* FUNGICIDE

ACTIVE CONSTITUENT: 375G/KG COPPER PRESENT AS COPPER OXYCHLORIDE

APPLICATION RATES

TREE AND VINE CROPS:

CROP	DISEASE	STATE	RATE
Apricots	Freckle (<i>Venturia carpophila</i>) Shot hole (<i>Stigmima carpophila</i>)	All states	335 g/100 L
Cherries, Peaches, Nectarines, Plums	Shot hole (<i>Stigmima carpophila</i>)		
Almonds		Qld, Vic, Tas, SA, WA only	
Apricots, Cherries	Bacteria gummosis (<i>Pseudomonas syringae</i>)	Qld, Vic, Tas, SA, WA only	335 g/100 L
		NSW, Vic, Tas, SA, WA only	200 g/100 L
Peaches, Nectarines		All states	335 g/100 L
Almonds	Leaf Curl (<i>Taphrina deformans</i>)	Qld, Vic, Tas, SA, WA only	
Stone Fruit	Blossom Blight (<i>Monolinia fructicola</i>) Freckle (<i>Venturia carpophila</i>) Rust (<i>Tranzschelia discolor</i>)	NSW, Vic, Tas, SA, WA only	535 g/100 L
	Leaf Curl (<i>Taphrina deformans</i>) Shot hole (<i>Stigmima carpophila</i>)		
	Bacterial Spot (<i>Xanthomonas arboricola</i>) Bacterial Canker (<i>Pseudomonas syringae</i>) Leaf Curl (<i>Taphrina deformans</i>) Shot hole (<i>Stigmima carpophila</i>)	Qld only	1000 g/100 L 665 g/100 L (Apricots)
Apples Pears	Black Spot (scab) (<i>Venturia inaequalis</i>) (<i>Venturia pirina</i>)	All states	335 g/100 L and winter oil
Avocado	Anthracoese (<i>Glomerella conglutata</i>)	NSW, Qld, SA, Vic, NT only	335 g/100 L
Citrus	Black Spot (<i>Phoma citricarpa</i>) Melanose (<i>Phoma citri</i>)	NSW, SA, Vic, WA only	235 g/100 L & summer oil
	Black Spot (<i>Phoma citricarpa</i>) Smoky Blotch (<i>Gloeodes pomigena</i>) Scab (lemons) (<i>Elsinoe fawcettii</i>)	Qld, NT only	265-400g /100 L plus 600 ml summer oil
Durians	Pink Disease, Anthracnose	All states	535 g/100 L
	Phytophthora Root Rot (<i>Phytophthora spp.</i>)	All states	535 g/100 L
Guavas	Anthracoese (<i>Colletotrichum psidii</i>) Algal Leaf Spot (<i>Cephaleuros virescens</i>)	NSW, NT only	335 g/100 L
Lychee	Parasitic algae (<i>Cephaleuros virescens</i>)	Qld, NSW only	535 g/100 L plus 20 ml of a 100% wetting agent
Macadamias	Husk Spot (<i>Cercoseptira spp.</i>)	Qld, NSW, NT only	335 g/100 L
	Anthracoese/Husk Rot (<i>Colletotrichum spp.</i>)	Qld, NSW, NT only	335 g/100 L
	Pink Limb Blight (<i>Corticium salmonicola</i>)		
Mangoes	Bacterial Black Spot (<i>Xanthomonas campestris</i>)	Qld, NSW, WA only	335 g/100 L
Mangosteens	Pink Disease, Anthracnose, Leaf Spots	All states	535 g/100 L
Passionfruit	Brown Spot (<i>Alternaria passiflorae</i>)	Qld only	265 g/100 L
	Septoria Spot (<i>Septoria passiflorae</i>)	NSW, NT, SA, Tas, Vic, WA only	400-535 g/100 L
	Phytophthora Blight (<i>Phytophthora nictianae</i>)	Qld, NSW only	135 g/100 L
Rambutans	Pink Disease, Anthracnose, Leaf Spots	All states	535 g/100 L
Vines	Downy Mildew (<i>Plasmopara viticola</i>)	NSW, Qld, SA, Tas, Vic, WA only	270 g/100 L
Walnuts	Walnut Blight (<i>Xanthomonas campestris</i>)	All states	400 g/100 L plus 175 ml summer oil

NON-TREE AND VINE CROPS:

CROP	DISEASE	STATE	RATE
Bananas	Cercospora, Leaf Spot (<i>Cercospora musae</i>)	NSW, Qld, WA only	335 g/100 L plus 600 ml summer oil/100 L or 3.4 kg/ha
Beans	Common Blight (<i>Xanthomonas campestris</i>)	SA, Tas, Vic, WA only	335 g/100 L or 3.4 kg/ha
	Bacterial Brown Spot (<i>Pseudomonas syringae pv syringae</i>)	All states	335 g/100 L or 3.4 kg/ha
	Halo Blight (<i>Pseudomonas syringae pv phaseolicola</i>)	SA, Tas, Vic, WA only	
		NSW, Qld, NT only	335-400 g/100L or 3.4-4 kg/ha
	Rust (<i>Uromyces spp.</i>) Chocolate Spot (<i>Botrytis spp.</i>)	SA, Vic, WA only	335 g/100 L or 3.4 kg/ha
Faba Beans	Rust (<i>Uromyces spp.</i>)	All states	335 g/100 L or 3.4 kg/ha
	Chocolate Spot (<i>Botrytis spp.</i>)	NSW, Qld, SA, Tas, WA only	
Brassicas	Black Spot (<i>Xanthomonas campestris</i>) Peppery Leaf Spot (<i>Pseudomonas syringae</i>) Ring Spot (<i>Mycosphaerella brassicicola</i>) Downy Mildew (<i>Peronospora parasitica</i>)	All states	335 g/100 L or 3.4 kg/ha
Capsicums	Bacterial Spot (<i>Xanthomonas vesicatoria</i>) Bacterial canker (<i>Corynebacterium michiganense</i>)	All states NSW, Qld, SA, Tas, WA only	335 g/100 L or 3.4 kg/ha
Carrots	Leaf Spot (<i>Alternaria spp.</i>)	All states	335 g/100 L or 3.4 kg/ha
Celery	Septoria, Leaf Spot (<i>Septoria apiicola</i>) Early Blight (<i>Cercospora apii</i>)	All states NSW, Vic, SA, WA, NT only	335 g/100 L or 3.4 kg/ha
Cucurbits i.e., Rockmelon, Watermelon, Cucumber	Angular Leaf Spot (<i>Pseudomonas syringae</i>) Bacterial Leaf Spot (<i>Xanthomonas campestris</i>)	All states	335 g/100 L or 3.4 kg/ha
Lettuce	Downy Mildew (<i>Bremia lactucae</i>) Bacterial Leaf Spot (<i>Xanthomonas campestris</i>) Anthracnose (<i>Marssonina panationiana</i>)	All states NSW, SA, Tas, Vic, WA only	335 g/100 L or 3.4 kg/ha
Onions	Downy Mildew (<i>Peronospora destructa</i>)	All states	335 g/100 L or 3.4 kg/ha
Ornamentals (flowers & shrubs)	Fungal Leaf Spots, Downy Mildew	All states	400 g/100 L or 4.0 kg/ha
Parsnips	Leaf Spots (<i>Septoria spp.</i>)	SA, Vic, WA only	335 g/100 L or 3.4 kg/ha
Peas	Ascochyta Blight (<i>Ascochyta spp.</i>) Bacterial Blight (<i>Pseudomonas syringae</i>)	All states	335 g/100 L or 3.4 kg/ha
Potatoes	Target Spot (<i>Alternaria solani</i>) Irish Blight (<i>Phytophthora infestans</i>)	All states	250-300 g/100L or 2.5-3.0 kg/ha
Red Beet	Downy Mildew (<i>Peronospora fasiona</i>) Rust (<i>Uromyces betae</i>)	All states	335 g/100 L or 3.4 kg/ha
Rhubarb	Crown Rot (<i>Phytophthora spp.</i>) Downy Mildew (<i>Peronospora jacciana</i>)	All states	335 g/100 L or 3.4 kg/ha
Silver Beet, Spinach	Downy Mildew (<i>Peronospora fasiona</i>)	All states	335 g/100 L or 3.4 kg/ha
Tomatoes	Bacterial Spot (<i>Xanthomonas campestris</i>) Bacterial Speck (<i>Pseudomonas syringae</i>) Bacterial Canker (<i>Corynebacterium michiganense</i>) Target Spot / Early Blight (<i>Alternaria solani</i>)	All states	295-335 g/100L or 2.5 kg/ha
	Septoria Leaf Spot (<i>Septoria spp.</i>) Irish Blight/ Late Blight (<i>Phytophthora infestans</i>)	All states	270-290 g/100L or 2.5 kg/ha



ORGANIC CROP PROTECTANTS
1956 Dandenong Rd, Clayton VIC 3168
Telephone: 1800 634 204 (Free Call)
ocp.com.au

