

Aminogro[®]

12% Copper

MAXi

Organic Carbon: 18% w/w Amine Nitrogen: 6% w/w

 **Organoplex[®] L-form Amino Acid**
Chelate Carboxylate Mannitol Complex

PLANT FUNCTION

- Photosynthesis
- Enzyme production
- Carbohydrate & nitrogen metabolism
- Lignin synthesis for cell wall strength and root cap

DEFICIENCY SYMPTOMS

- Dieback of stems & twigs
- Yellowing of leaves
- Stunted growth



ORGANIC CROP PROTECTANTS
1956 Dandenong Rd, Clayton VIC 3168
Telephone: 1800 634 204 (Free Call)
ocp.com.au


**Input for
Organic
Production**
ACO CERT. NO. 222
Restricted for use under ACO Standards

CONTENTS
25 kg

GENERAL INFORMATION

Aminogro® MAXi Copper provides plants with a copper chelate using L-form free amino acids and other simple organic sugars and acids. Nutrients are absorbed and shuttled around inside plants in the form of amino acids, simple sugars and acids so OrganoPLEX™ is an efficient way of delivering and mobilising copper in plants. The organic carbon in Aminogro® MAXi Copper also enhances microbial activity in soil which assists the plant accessing copper and other nutrients when applied to the soil.

DIRECTIONS FOR USE:

The rates below are a guide and dependent upon soil and sap tests throughout the growing season. Consult your OCP Agronomist for further advice.

Tree Crop	Rate	Application Timing
Pome Fruit	800 g-1.5 kg/ha	1-2 weeks before flowering
Stone Fruit	800 g-1.5 kg/ha	At first flush
Nuts	800 g-1.5 kg/ha	At first flush
Bananas	500 g-1 kg/ha	With seasonal application of potassium and nitrogen
Vine Crop	Rate	Application Timing
Table Grapes	500 g-1.5 kg/ha	2 weeks before flowering
Wine Grapes	800 g-1.5 kg/ha	2 weeks before flowering
Strawberries/ Blueberries	500 g-1 kg/ha	Begin just before flowering & repeat as sap results indicate
Protected Crops	Rate	Application Timing
Hydroponic run to waste & NFT Crops	100 g/10,000 L	Consult OCP agronomist for crop specific requirements
Nursery Stock	Foliar	If deficiency is detected consult OCP agronomist

Field Crops	Rate	Application Timing
Cereal, Soybeans	1 kg/ha	As required by soil test results
Leafy veggies, Lettuce	500 g/ha	4 leaf stage
Carrots, Onions, Leeks, Turnips	500 g/ha	2 weeks after emergence
Brassica, Crucifer	500 g/ha	6-8 true leaf stage
Bean & Peas	500 g/ha	4-6 true leaf stage
Potatoes, Yam, Taro	500 g/ha	At tuber set
Sweet Corn	500 g/ha	2-6 true leaf stage
Tomato, Capsicum	500-750 g/ha	15-20cm
Cucurbits	500 g/ha	4 true leaf stage
Professional Turf	Rate	Application Timing
Greens & Tees	100 g/1000 m ²	As required by soil test. Higher rates may be used during time of disease pressure
Fairways	500 g/ha	As required by soil test

COMPATIBILITY

Aminogro® MAXi Copper can be mixed with most horticultural sprays. However, the performance of some sprays may be affected by high cation levels like calcium, potassium and phosphorus. It is best to apply these sprays separately or conduct a jar test to determine compatibility in the ratios they are to be used. Always check pesticide labels for compatibility with foliar fertilisers, or test the mixture on a small area before widespread application.

For further information contact Organic Crop Protectants

Free Call: 1800 634 204 Email: info@ocp.com.au

FOR EMERGENCIES ONLY, CALL AUSTRALIA 1800 033 111 (ALL HOURS)

Aminogro and the OrganoPLEX device are registered trade marks and the OCP drop device is a trade mark of Organic Crop Protectants

