

Aminogro[®]

12% Iron

MAXi

Organic Carbon: 19% w/w Amine Nitrogen: 7.6% w/w

 **Organoplex[®]** L-form Amino Acid
Chelate Carboxylate Mannitol Complex

PLANT FUNCTION

- Chlorophyll production
- Photosynthesis
- Nitrogen uptake & lignin formation

DEFICIENCY SYMPTOMS

- Yellowing of leaves and interveinal chlorosis (green veins and yellowing between veins)



ORGANIC CROP PROTECTANTS

1956 Dandenong Rd, Clayton VIC 3168
Telephone: 1800 634 204 (Free Call)
ocp.com.au



CONTENTS
25 kg

GENERAL INFORMATION

Aminogro® MAXi Iron provides plants with an iron chelate using L-form free amino acids and other simple organic sugars and acids. Nutrients are absorbed and shuttled around inside plants in the form of amino acids, simple sugars and acids so OrganoPLEX™ is an efficient way of delivering and mobilising iron in plants. The organic carbon in Aminogro® MAXi Iron also enhances microbial activity in soil which assists the plant accessing iron and other nutrients when applied to the soil.

DIRECTIONS FOR USE:

The rates below are a guide and dependent upon soil and sap tests throughout the growing season. Consult your OCP Agronomist for further advice.

| Tree Crop | Rate | Application Timing |
|------------------------------|-----------------|---|
| Pome Fruit | 1-2 kg/ha | 2 weeks after fruit set |
| Stone Fruit | 1-2 kg/ha | 1- 2 weeks after fruit set |
| Nuts | 1-2 kg/ha | 2 -3 weeks after nut set |
| Vine Crop | Rate | Application Timing |
| Table Grapes | 1-1.5 kg/ha | Pre-bloom E-L 12 & as required by sap & soil test |
| Wine Grapes | 750 g-1.5 kg/ha | Pre-bloom E-L 12 & as required by sap test |
| Strawberries/ Blueberries | 750 g-1.5 kg/ha | Before flowering & as required by sap & soil test |
| Protected Crops | Rate | Application Timing |
| Hydroponic NFT All Crops | 200 g/10,000 L | Consult OCP agronomist |
| Nursery Stock | Foliar | If deficiency is detected consult OCP agronomist |

| Field Crops | Rate | Application Timing |
|------------------------------------|---------------------------|---|
| Cereal, Soybeans | 2-3 kg/ha | As required by soil test results |
| Leafy Veggies Lettuce | 1-2 kg/ha | 4 leaf stage |
| Carrots, Onions, Leeks, Turnips | 2 kg/ha | 2 weeks after emergence |
| Brassica, Crucifer | 2 kg/ha | 6-8 true leaf stage |
| Bean & Peas | 2 kg/ha | 4-6 true leaf stage |
| Potatoes, Yam, Taro | 2 kg/ha | At tuber set |
| Sweet Corn | 2 kg/ha | 1. 2-6 true leaf stage 2. 50-75cm growth |
| Cucurbits | 2 kg/ha | 4 true leaf stage |
| Tomato, Capsicum | 750 g-1 kg/ha | 15-20cm |
| Professional Turf | Rate | Application Timing |
| Greens & Tees | 500 g/1000 m ² | As required for colour |
| Fairways | 2 kg/ha | As required for colour |

COMPATIBILITY

Aminogro® MAXi Iron can be mixed with most horticultural sprays. However, the performance of some sprays may be affected by high cation levels like calcium, potassium and phosphorus. It is best to apply these sprays separately or conduct a jar test to determine compatibility in the ratios they are to be used. Always check pesticide labels for compatibility with foliar fertilisers, or test the mixture on a small area before widespread application.

For further information contact Organic Crop Protectants

Free Call: 1800 634 204 Email: info@ocp.com.au

FOR EMERGENCIES ONLY, CALL AUSTRALIA 1800 033 111 (ALL HOURS)

Aminogro and the OrganoPLEX device are registered trade marks and the OCP drop device is a trade mark of Organic Crop Protectants

