



Organic Crop Protectants

READ SAFETY DIRECTIONS
BEFORE OPENING OR USING

HELICOVEX[®]

BIOLOGICAL INSECTICIDE

ACTIVE CONSTITUENT: 7.5×10^9 POLYHEDRAL INCLUSION BODIES OF THE
NUCLEOPOLYHEDROVIRUS OF *Helicoverpa armigera* PER MILLILITRE
FOR THE CONTROL OF *Helicoverpa* spp LARVAE IN VARIOUS CROPS AS
SPECIFIED IN DIRECTIONS FOR USE

IMPORTANT: READ PRODUCT INFORMATION UNDER LABEL BEFORE USE



80%
Magnification
Barcode
9336099556755

Y20-3241

CONTENTS
5L



ORGANIC CROP PROTECTANTS
1956 Dandenong Road, Clayton VIC 3168
Telephone: 1800 634 204 (Free Call)
www.ocp.com.au

GENERAL INSTRUCTIONS

HELICOVEK (nucleopolyhedrovirus) is a highly specific naturally occurring pathogen of *Helicoverpa* spp. The effectiveness of HELICOVEK is dependent on a number of important factors: environmental conditions, application and the feeding behaviour of the pest. It is because of the requirement for near perfect conditions that the performance of HELICOVEK is variable and at times, the level of control may be below expectations. The speed of activity of HELICOVEK is also dependent on climatic conditions. Larvae can take up to 8 days to die. Daytime temperatures of 25°C to 35°C are ideal for the activity of HELICOVEK.

Good coverage of the feeding sites of the larvae is essential, as the product needs to be ingested to be effective. HELICOVEK will not control larvae that do not feed on treated areas, e.g. when larvae are feeding in protected feeding sites such as inside cotton bolls, lettuce hearts, bean pods, corn cobs and flowers.

Temperature and Larval Size Spray Decision Table

Larval Size	Temperature (°C)			
	<18	18-25	25-35*	>35
0-3 mm	X	33	33	X
4-7 mm	X	3	33	X
8-13 mm	X	3	3	X
>13 mm	X	X	X	X

33 Ideal Conditions; 3 Good Conditions; X Consider options

* Avoid spraying or increase application volume when temperature is above 30°C and humidity is below 40%.

Good coverage is the key factor in ensuring maximum performance of HELICOVEK. For this reason, apply HELICOVEK to coincide with optimum environmental conditions, such as periods of high humidity that can occur in the early hours of the morning (2 to 8 am). This should still coincide with warm (>25°C) conditions to ensure high larval activity. If HELICOVEK is applied during hot and dry conditions, increase application volumes and droplet size to improve the level of coverage achieved.

MIXING

Shake the container well before use. **Spray water pH should be between 5 to 8.5** - spray water pH above 9 will damage the virus and performance will be reduced. If needed, use a suitable buffer or acidifier. Partially fill the spray tank with water, add the required amount of HELICOVEK while agitating and top up the spray tank with water to the required volume. HELICOVEK should be applied as soon after mixing as possible. The virus can be rendered inactive if the mixture is left to stand overnight.

APPLICATION

Use application parameters (nozzles, swath width, pressure, boom height, speed, etc.) to ensure thorough coverage of the target area.

Horticultural crops:

Apply by ground rig or hand held equipment in a minimum of 400 litres of water per hectare.

Broadacre crops:

Ground Rig: Apply in a minimum of 100 litres of water per hectare.

Aerial - High Volume: Apply in a minimum of 30 litres of water per hectare. This application method is particularly susceptible to droplet evaporation, especially during hot and dry conditions (temperature greater than 30°C and humidity less than 40%). Droplet evaporation will reduce coverage, which can have a detrimental impact on performance. During hot and dry conditions avoid using this application method - wait until conditions favour good coverage or apply in ULV (see below). Alternatively, if application in water by air during hot and dry conditions cannot be avoided, increase application volume and/or use an anti-evaporation additive to improve coverage.

Aerial - Ultra-Low Volume (ULV):

Use an approved carrier such as Synertrac Horti Oil at 1-2 Litres per hectare; or D-C-Tron, Canopy or Biopest Oil applied in a minimum volume of 3 litres per hectare. When applying HELICOVEK in ULV, do not tank mix with other pesticides or fertilisers (refer to Compatibility).

Via Overhead Irrigation:

HELICOVEK can be effectively applied to crops in overhead irrigation water. The product should be introduced to the irrigation water at the appropriate rate using filtration equipment. If the product is diluted in water prior to injection into the irrigation water, ensure that the dilution water is clean and not silty with a pH of 7 or less and ensure there is constant agitation. Preferably, rainwater should be used for dilution. Ensure any diluted HELICOVEK is used within 10 hours of mixing. For one-pass mobile irrigators such as centre pivots and laterals, continuously introduce the required quantity of HELICOVEK into the irrigation water over the course of irrigation. Apply HELICOVEK in no more than 10 mm of irrigation water. For static irrigators, introduce the required amount of HELICOVEK into the irrigation water just prior to completion of the irrigation period, to maximise the concentration of HELICOVEK applied and the amount that remains on the crop.

COMPATIBILITY

In water HELICOVEK is highly compatible with the majority of pesticides and fertilisers when mixed in water. Ensure that the mixture has a pH between 5 and 8.5 before adding HELICOVEK as alkaline pH (greater than 9) will damage the virus.

In ULV For ULV application in oil, HELICOVEK is not compatible with other pesticides because the undiluted solvents in these products can damage the virus.

PRECAUTIONS

Re-entry: Do not allow entry into treated areas until spray has dried. When prior entry is necessary,

wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

Flaggers: Do not use human flaggers/markers unless they are protected by engineering controls such as enclosed cabs.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Do not contaminate streams, rivers or waterways with the product, including via run-off, spray drift or disposal of used containers.

STORAGE AND DISPOSAL

Storage: Keep out of reach of children. Store in the original container out of direct sunlight or at below 4°C. The product is stable for 2½ years from DOM if stored as indicated.

Disposal: Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted product on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin and open wounds. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. When opening the container, prepare spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow length PVC gloves and a face shield or goggles. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Ph. 131 126). FOR EMERGENCIES ONLY, CALL: AUSTRALIA 1800 033 111.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which can be obtained from the supplier and ocp.com.

HELICOVEK is a registered trade mark of Andermatt Biocontrol AG. The OCP Drop Device is a trade mark of Organic Crop Protectants. APVNA Approval Number: 68347/100611

HELICOVEX® BIOLOGICAL INSECTICIDE

ACTIVE CONSTITUENT: 7.5 X 10⁹ POLYHEDRAL INCLUSION BODIES OF THE NUCLEOPOLYHEDROVIRUS OF *Helicoverpa armigera* PER MILLILITRE

DIRECTIONS FOR USE

Crop/Situation	Pest	Rate (L/ha)	Critical Comments
Cereal grains including: Maize, Popcorn, Sorghum Lucerne (Alfalfa)	<i>Helicoverpa armigera</i>	100 mL/ha	All Crops: Thorough coverage of the crop is essential as HELICOVEX needs to be ingested to be effective. Refer to the General Instructions and Application sections for additional application advice. HELICOVEX should be applied between 25°C and 35°C when larvae are actively feeding. HELICOVEX is more effective on smaller larvae. Target application to coincide with neonate larvae emerging from their eggs. HELICOVEX should not be applied on larvae larger than 13 mm in length. HELICOVEX will provide between 60 and 90% control. Under extremely high pest pressure or sub-optimal application conditions, or when protection against damage is vital, additional control options should be considered. Sorghum: Application should be made 3 days after 50% of panicles have reached 100% flowering. Linseed: Use a non-ionic surfactant at the manufacturer's specified rate to improve coverage.
Oilseed including: Canola, Linseed, Mustard seeds, Peanut, Safflower, Sesame seed, Sunflower Potatoes Pulses including: Azuki bean, Broad bean, Chickpea, Cowpea, Faba bean, Field pea, Kidney bean, Lablab, Lentil, Lima bean, Lupin, Mung bean, Navy bean, Pigeon pea, Soybean, Vetch	Corn earworm/ Cotton bollworm/ Tobacco budworm AND <i>Helicoverpa punctigera</i> Native budworm	130 mL/ha OR 130 mL/ha + a registered larvicide at its label rate	
Cotton		130 mL/ha	Chickpeas: The addition of powdered milk at a rate of 1.0 kg/ha may improve the performance of HELICOVEX in this crop. HELICOVEX is unlikely to reduce larval numbers below threshold if the initial population exceeds 6 per metre of row. Cotton: HELICOVEX should not be applied on larvae larger than 7 mm in length. When applied alone, HELICOVEX is unlikely to reduce larval numbers below threshold if the initial population exceeds 4 per metre of row. HELICOVEX should be used in accordance with the Cotton Best Management Practices Manual.
Sweetcorn		130 mL/ha	Sweetcorn: Application should be made from the early vegetative growth stage through to tasselling and prior to the emergence of silks. HELICOVEX has short residual activity and re-treatment may be required at 2 to 3 day intervals, depending on egg counts and crop growth rates.
Berryfruit including: Blackberries, Blueberries, Boysenberry, Cranberry, Currants, Gooseberry, Raspberries, Strawberry Brassica vegetables including: Brassica leafy vegetables, Broccoli, Brussels sprouts, Cabbages, Cauliflower, Chinese broccoli Celery Cucurbits including: Cucumber, Melons, Pumpkins, Squash, Watermelon, Zucchini Fruiting vegetables including: Eggplant, Peppers (capsicum and chilli), Tomato Leafy vegetables including: Endive, Lettuce, Rockette (Rucola), Silver beet, Spinach Legume Vegetables including: Green beans, Green peas, Snow peas, Sugar snap peas Ornamental flowers and plants Pome fruit including: Apples, Nashi, Pears		100 to 200 mL/ha	Horticultural crops: Use a higher rate when flowers, fruit or economic parts of the crop are present, under high pest pressure conditions or to control larvae greater than 7 mm in length. Use lower rates during vegetative stages of crop production. HELICOVEX has a short residual activity and re-treatment may be required at 2 to 3 day intervals. Use a non-ionic surfactant at the manufacturer's specified rate to improve coverage.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

STORAGE AND DISPOSAL

Storage: Keep out of reach of children. Store in the closed, original container out of direct sunlight at or below 4°C. The product is stable for 2½ years from DOM if stored as indicated.

Disposal: Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted product on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin and open wounds. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. When opening the container, preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow length PVC gloves and a face shield or goggles. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Ph. 131126) FOR EMERGENCIES ONLY, CALL: AUSTRALIA 1800 033 111.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which can be obtained from the supplier and ocp.com.au.

ORGANIC CROP PROTECTANTS 1956 Dandenong Road, Clayton VIC 3168.

Telephone: 1800 634 204 (Free Call) ocp.com.au

APVMA Approval Number: 68347/100611