



Typical Analysis	% w/w
Nitrogen (N) as Nitrate	5.2
Nitrogen (N) as Ammonium	7.8
Total Nitrogen	13.0
Phosphorus (P) as water soluble	3.9
Total Phosphorus	4.8
Total Potassium (K) as sulfate	9.1
Sulfur (S) as Sulfates	5.0
Magnesium (Mg) as Sulfate	1.2
Iron (Fe) as EDTA chelate	0.2
Manganese (Mn) as Sulfate	0.06
Copper (Cu) as Sulfate	0.05
Zinc (Zn) as Sulfate	0.015
Boron (B) as Boric Acid	0.02
Molybdenum (Mo) as Sodium Molybdate	0.02

Application Precautions

TRIAL PRIOR TO FERTILISER CHANGE: We recommend that this product is trialled under local growing conditions prior to use on selected crops so that application methods and desired rates can be validated.

- Not suitable for use on species which are highly sensitive to phosphorus. For these species we recommend Nutricote Standard Purple (100 day), Pink (180 day) and N18.7 Total range.
- Avoid application to plants under stress or seek advice on reduced rates.
- Avoid mounding fertiliser against the base of the plant.
- Plant nutrients can cause staining of porous stone and concrete surfaces.
- Keep away from pools, ponds and other bodies of water.
- If incorporating into potting mix use within 14 days and if temperatures are in excess of 30°C irrigate thoroughly after potting to leach potential excess fertiliser from mix.
- Potting as soon as possible after incorporation in mixes is preferred to ensure longevity of fertiliser release and cost efficiency of fertiliser program.
- If using in potting mixes with high cation exchange capacities use lower recommended rates or seek advice prior to use.
- Always water in well after application.

Caution

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed

Open bags must be resealed after use by tying before storage, especially in winter and in areas of high humidity. Failure to do so may impact on future product performance.

The use of this product is beyond the control of the seller, no condition or warranty is given and none shall be implied as to its suitability and effectiveness for any particular purpose. The seller accepts no responsibility for any consequences whatsoever from the use of the product.

Yates Australia
1 Gow Street, Padstow
NSW, 2211 Australia

Sales Support: 1300 132 153

www.yatescommercial.com.au

© Yates and The Squares Device are registered trade marks.

© Nutricote is a registered trade mark of JCAM Agri. Co. Ltd, Tokyo Japan.

The front panel of this package is owned under copyright laws by Yates Australia.

25kg

Y 39-352



Controlled Release Fertiliser
Nutricote
N13 Total+TE 25kg




N13 Total+TE 25kg



Controlled Release Fertiliser
Nutricote

140 DAY



Controlled Release Fertiliser
Nutricote


N13 Total+TE

- Medium term feeding for flowering trees and shrubs
- Enhances flowering and root growth
- Trace elements in every prill

140 DAY

Soil temperature	10°C	15°C	20°C	25°C	30°C	35°C
Release time (days)	280	220	180	140	115	90

25kg



Total+TE


N 13.0

P 4.8

K 9.1



5H4/Y25.2/S/11/AUS/YATES.30747



9 310428 009051

140 DAY

Controlled Release Fertiliser
Nutricote



N13 Total+TE 25kg



AMMONIUM NITRATE
BASED FERTILISER
UN 2071



MEASUREMENTS (spoons and cups)

1 teaspoon (tsp) = 5g

1 tablespoon (tbs) = 15g

1/2 cup = 135g

2 cups = 540g

Controlled Release Fertiliser
Nutricote

N13 Total+TE 140 Day

Benefits

- Medium term feeding for flowering trees and shrubs.
- Enhances flowering and root growth.
- Trace elements in every prill.
- Designed for hot fluctuating temperatures.
- Superior control of nutrient release over 140 days @ 25°C.
- Reduced leaching so optimum fertiliser levels are maintained.
- A durable, resilient and uniform polymer coating on each prill.
- Every prill is homogeneous to ensure consistent analysis.
- Consistent prill size for easier and more precise mechanical application.
- Consistent rate of nutrient release providing steady feeding according to plant nutritional needs.

How to apply

Top dressing

- Evenly apply to the base of the plant keeping fertiliser away from the main trunk or stem.
- Water in well after application with a rose type spray to minimise water blast of prills from pot.

Incorporation

- Mix well into the soil profile ensuring even application.

General

- Release period is governed by soil temperature as shown below.
- Nutricote should provide the entire requirements of nitrogen, potassium and maintenance levels of phosphorus and trace elements for the release period.
- Use high rates for fast draining soils or mixes of low fertility and for species that are fast growing and watered frequently.
- Use medium rates when:
 - a) using on sensitive crops.
 - b) soil temperatures are very high.
 - c) liquid feeding two to three times per week.

Release rates

Soil temperature	10°C	15°C	20°C	25°C	30°C	35°C
Release time (days)	280	220	180	140	115	90

Suggested application rates

Top dressing (g)

Pot size (diameter)	Medium (grams per pot)	High (grams per pot)
100mm (4")	2	3
125mm (5")	4	6
150mm (6")	6	8
175mm (7")	9	15
200mm (8")	11	18
250mm (10")	18	30
300mm (12")	24	40
Incorporation (kg/m²)	4	5-6



Controlled Release Fertiliser
Nutricote

N13 Total+TE 25kg

140 DAY



9 310428 009051

