

Typical Analysis	% w/w
Nitrogen (N) as Nitrate	5.5
Nitrogen (N) as Ammonium	8.5
Total Nitrogen (N)	14.0
Phosphorus (P) as water soluble	4.8
Total Phosphorus (P)	5.7
Total Potassium (K) as sulfate	10.8
Sulfur (S) as Sulfates	4.0

Application Precautions

TRIAL PRIOR TO FERTILISER CHANGE: We recommend that this product is trialled under local growing conditions prior to use on selected crops so that application methods and desired rates can be validated.

- Not suitable for use on species which are highly sensitive to phosphorus. For these species we recommend **Nutricote** Standard Purple (100 day), Pink (180 day) and N18.7 Total range.
- Avoid application to plants under stress or seek advice on reduced rates.
- Avoid mounding fertiliser against the base of the plant.
- Plant nutrients can cause staining of porous stone and concrete surfaces.
- Keep away from pools, ponds and other bodies of water.
- If incorporating into potting mix use within 14 days and if temperatures are in excess of 30°C irrigate thoroughly after potting to leach potential excess fertiliser from mix.
- Potting as soon as possible after incorporation in mixes is preferred to ensure longevity
- of fertiliser release and cost efficiency of fertiliser program.
- If using in potting mixes with high cation exchange capacities use lower recommended rates or seek advice prior to use.
- Always water in well after application.

Caution

KEEP OUT OF REACH OF CHILDREN

Open bags must be resealed after use by tying before storage, especially in winter and in areas of high humidity. Failure to do so may impact on future product performance.

The use of this product is beyond the control of the seller, no condition or warranty is given and none shall be implied as to its suitability and effectiveness for any particular purpose. The seller accepts no responsibility for any consequences whatsoever from the

Important Product Disclaimer

Yates makes no representations or warranties that this product will be suitable for use. Subject to any applicable legislation which cannot be excluded (such as the any failure in performance of this product or any losses, damages, or injuries (consequential or otherwise) arising from the storage, mixing, application or other use of this product or for any representations made by you about the product which have not been approved in writing by **Yates**. **Yates** recommends that you contact an agronomist or suitably qualified person prior to the application of this product and/or conduct trials of this product prior to full commercial application.

1 Gow Street, Padstow NSW, 2211 Australia

Sales Support: 1300 132 153 www.yatescommercial.com.au ® Yates and The Squares Device are registered trade marks of DuluxGroup (Australia) Pty Ltd. ® Nutricote is a registered trade mark of JCAM Agri. Co. Ltd, Tokyo Japan. The front panel of this package is owned under copyright laws by Yates Australia.

Y 39-350



Standard Silver 25kg






Standard ates

Silver

T





Yates®



360

DAY

Controlled Release Fertiliser

Nutre

Standard Silver

- Long term feeding for super advanced fruiting and flowering plants
- Once a year fertiliser for large trees and shrubs
- Robust coating for increased safety

Soil temperature	10°C	15°C	20°C	25°C	30°C	35°C
Release time (days)	920	740	535	360	250	165

25kg



Standard



P 5.7

K 10.8



(N) 5H4/Y25.2/S/13/AUS/YATES.30747

Controlled Release Fertiliser Nutricate

Standard Silver 360 Day

Benefits

- Long term feeding for super advanced fruiting and flowering plants.
- Once a year fertiliser for large trees and shrubs.
- Robust coating for increased safety.
- Designed for hot fluctuating temperatures.
- Superior control of nutrient release over 360 days @ 25°C.
- Reduced leaching so optimum fertiliser levels are maintained
- A durable, resilient and uniform polymer coating on each prill.
- Every prill is homogeneous to ensure consistent analysis.
- Consistent prill size for easier and more precise mechanical application.
- Consistent rate of nutrient release providing steady feeding according to plant nutritional needs.

How to apply

Top dressing

Evenly apply to the base of the plant keeping fertiliser away from the main trunk or stem.

• Water in well after application with a rose type spray to minimise water blast of prills from pot.

Incorporation

Mix well into the soil profile ensuring even application.

General

C3te®

Controlled Release Fertiliser

Yates

Standard

- Release period is governed by soil temperature as shown below
- Nutricote should provide the entire requirements of nitrogen, potassium and maintenance levels of phosphorus for the release period
- Use high rates for fast draining soils or mixes of low fertility and
- for species that are fast growing and watered frequently.
- Use medium rates when:
- a) using on sensitive crops.
- b) soil temperatures are very high.
- c) liquid feeding two to three times per week.

Release rates

Soil temperature	10°C	15°C	20°C	25°C	30°C	35°C
Release time (days)	920	740	535	360	250	165

MEASUREMENTS


= 15g

(spoons and cups)

Suggested application rates

Top dressing (q)

Pot size (diameter)	Medium (grams per pot)	High (grams per po
100mm (4")	3.5	5
125mm (5")	7	10
150mm (6")	8	12
175mm (7")	12	20
200mm (8")	18	30
250mm (10")	30	50
300mm (12")	42	70
Incorporation (kg/m	³) 6–7	8-9









Controlled Release Fertiliser

Nutricate **Standard Silver 25kg**

360 DAY

