



Controlled Release Fertiliser Nutricote®

Standard Gold


- Medium-long term feeding for foliage plants
- Suitable for most natives and exotics
- Ideal for top dressing semi advanced to super advanced plants



Growing media temperature	10°C	15°C	20°C	25°C	30°C	35°C
Release time (days)	610	380	265	180	135	95

Standard

N	16.0
P	4.4
K	8.3




Controlled Release Fertiliser
Nutricote®



Standard Gold 25kg

Standard Gold 25kg
Yates
Controlled Release Fertiliser
Nutricote®



25kg



Yates
Controlled Release Fertiliser
Nutricote®

Standard Gold 25kg



9 310428 572920



Typical Analysis	% w/w
Nitrogen (N) as Nitrate	6.8
Nitrogen (N) as Ammonium	9.2
Total Nitrogen	16.0
Phosphorus (P) as water soluble	2.2
Total Phosphorus	4.4
Total Potassium as sulfate	8.3
Total Sulfur (S) as sulfate	4.0

Application Precautions

TRIAL PRIOR TO FERTILISER CHANGE: We recommend that this product is trialled under local growing conditions prior to use on selected crops so that application methods and desired rates can be validated.

- Not suitable for use on species which are sensitive to phosphorus. For these species we recommend Nutricote Standard Purple (100 day), Pink (180 day) and the N18.7 Total range.
- Avoid application to plants under stress or seek advice on reduced rates.
- Avoid mounding fertiliser against the main trunk or stem of the plant.
- Plant nutrients can cause staining of porous stone and concrete surfaces.
- Keep away from pools, ponds and other bodies of water.
- If incorporating into potting mix use within 14 days and if temperatures are in excess of 30°C irrigate thoroughly after potting to leach potential excess fertiliser from mix.
- Potting as soon as possible after incorporation in mixes is preferred to ensure longevity of fertiliser release and cost efficiency of fertiliser program.
- If using in potting mixes with high cation exchange capacities use lower recommended rates or seek advice prior to use.
- Always water in well after application.

Caution

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid inhalation of dust, eye and skin contact. Wash hands and exposed skin after use.

FOR EMERGENCIES ONLY, CALL AUSTRALIA 1800 220 770 (ALL HOURS)

SDS: Additional information is listed in the Safety Data Sheet available from the supplier and ocp.com.au.

Storage: Always store in a cool, dry place out of direct sunlight. Open bags must be resealed after use by tying before storage, especially in winter and in areas of high humidity. Failure to do so may impact on future product performance.

Yates Australia
1956 Dandenong Rd
CLAYTON VIC 3168

Tel: 1800 634 204 (Free Call)
ocp.com.au

® Yates is a registered trade mark of DuluxGroup (Australia) Pty Ltd.
® Nutricote is a registered trade mark of JCAM Agri. Co., Ltd Japan.

25kg

Y35-1290



Controlled Release Fertiliser

Nutricote®

Standard Gold 180 Day

Benefits

- Superior control of nutrient release over 180 days @ 25°C.
- Medium-long term feeding for foliage plants.
- Suitable for most natives and exotics.
- Ideal for top dressing semi advanced to super advanced plants.
- Designed for hot fluctuating soil temperatures up to 35°C.
- Reduced leaching so optimum fertiliser levels are maintained.
- A durable, resilient and uniform resin coating on each prill.
- Every prill is homogeneous to ensure consistent analysis.
- Consistent prill size for easier and more precise mechanical application.
- Consistent rate of nutrient release providing steady feeding according to plant nutritional needs.

How to apply

Top dressing

- Evenly apply to the base of the plant keeping fertiliser away from the main trunk or stem.
- Water in well after application with a rose type spray to minimise water blast of prills from pot.

Incorporation

- Mix well into growing media ensuring even application.

General

- Release period is governed by growing media temperature as shown below.
- Nutricote should provide the entire requirements of nitrogen, potassium and maintenance levels of phosphorus for the release period.
- Use high rates for fast draining growing media of low fertility and for species that are fast growing and watered frequently.
- Use medium rates when:
 - a) using on sensitive crops.
 - b) growing media temperatures are very high.
 - c) liquid feeding two to three times per week.

Release rates

Growing media temperature	10°C	15°C	20°C	25°C	30°C	35°C
Release time (days)	610	380	265	180	135	95

Suggested application rates

Top dressing (g)

Pot Volume (Litres)	Pot Diameter	5 g/L Medium Rate (grams per pot)	6 g/L High Rate (grams per pot)
0.5	100mm (4")	2.5	3
1	125mm (5")	5	6
1.8	150mm (6")	9	11
3.0	175mm (7")	15	18
4.5	200mm (8")	23	27
8.0	250mm (10")	40	48
14	300mm (12")	70	84
Incorporation (kg/m ³)		5	6

MEASUREMENTS (spoons and cups)

- 1 teaspoon (tsp) = 5g
- 1 tablespoon (tbs) = 15g
- 1 cup = 270g



Controlled Release Fertiliser Nutricote®

Standard Gold 25kg

180
DAY

